

1 **Title:** Development and evaluation of novel real-time RT-PCR assays with locked nucleic acid
2 probes targeting the leader sequences of human pathogenic coronaviruses

3
4 **Running title:** Novel RT-PCR assays targeting CoV leader sequences

5
6 **Authors:** Jasper Fuk-Woo Chan,¹⁻⁴ Garnet Kwan-Yue Choi,² Alan Ka-Lun Tsang,² Kah-Meng
7 Tee,² Ho-Yin Lam,⁵ Cyril Chik-Yan Yip,² Kelvin Kai-Wang To,¹⁻⁴ Vincent Chi-Chung Cheng,²
8 Man-Lung Yeung,² Susanna Kar-Pui Lau,¹⁻⁴ Patrick Chiu-Yat Woo,¹⁻⁴ Kwok-Hung Chan,² Bone
9 Siu-Fai Tang,⁵ and Kwok-Yung Yuen^{1-4*}

10
11 **Affiliations:**

12 ¹State Key Laboratory of Emerging Infectious Diseases, ²Department of Microbiology,
13 ³Research Centre of Infection and Immunology, and ⁴Carol Yu Centre for Infection, The
14 University of Hong Kong; and ⁵Department of Pathology, Hong Kong Sanatorium and Hospital,
15 Hong Kong Special Administrative Region, China.

16
17 ***Correspondence:** Mailing address: Carol Yu Centre for Infection, Department of Microbiology,
18 The University of Hong Kong, Queen Mary Hospital, 102 Pokfulam Road, Pokfulam, Hong
19 Kong Special Administrative Region, China. Phone: (852) 22554892. Fax: (852) 28551241. E-
20 mail: kyyuen@hku.hk

21 **Keywords:** coronavirus, MERS, leader sequence, locked nucleic acid, LNA, PCR.

22 **Word count:** abstract = 50, text = 1837

23

24 **ABSTRACT**

25 Based on small-RNA-Seq analysis data, we developed highly sensitive and specific real-time
26 RT-PCR assays with locked nucleic acid probes targeting the abundantly expressed leader
27 sequences of MERS-CoV and other human coronaviruses. Analytical and clinical evaluations
28 showed their non-inferiority to a commercial multiplex PCR test for the detection of these
29 coronaviruses.

30 Coronaviruses (CoVs) have repeatedly crossed species barriers and some have emerged as
31 important human pathogens (1,2). HCoV-229E, HCoV-OC43, HCoV-NL63, and HCoV-HKU1
32 predominantly cause mild upper respiratory tract infections, while severe acute respiratory
33 syndrome CoV (SARS-CoV) and Middle East respiratory syndrome CoV (MERS-CoV)
34 frequently cause severe pneumonia with extrapulmonary manifestations (3-6). Highly sensitive
35 and specific laboratory diagnostic tests are essential for the control of emerging CoV outbreaks
36 (7). The gold standard of laboratory diagnosis of CoV infection is the isolation of infectious virus
37 from respiratory tract and/or other clinical specimens. However, most CoVs are either difficult or
38 dangerous to culture in cell lines (8,9). The need of convalescent samples and potential false-
39 positive results due to cross-reactivity with other CoVs limit the use of serum antibody detection
40 assays in the acute setting (10). The overall sensitivity of antigen detection assays is inferior to
41 that of molecular assays such as reverse transcription (RT)-PCR (11,12). With the increasing
42 availability of molecular diagnostic facilities and expertise in clinical microbiology laboratories
43 worldwide, RT-PCR has become the test of choice for diagnosing CoV infections (7,13-15).

44 Traditionally, the preferred targets of RT-PCR assays are genes that are conserved and/or
45 abundantly expressed from the viral genome (16). For CoVs, the most commonly employed
46 targets include the structural nucleocapsid (N) and spike (S) genes, and the non-structural RNA-
47 dependent RNA polymerase (RdRp) and replicase ORF1a/b genes (4,7). Recently, other unique
48 non-coding genome regions not present in related CoVs have also been utilized to develop RT-
49 PCR for the emerging MERS-CoV (7,13-15). The World Health Organization (WHO)
50 recommends using the upE assay (regions upstream of the envelope [E] gene) for laboratory
51 screening of suspected MERS cases, followed by confirmation with either the ORF1a or ORF1b
52 assays (7). Notably, a number of single nucleotide mismatches at different positions included in

53 the upE assay forward primer and probe have been detected in recent strains of MERS-CoV and
54 may affect the sensitivity of this assay (17). We hypothesize that additional gene targets may be
55 suitable for RT-PCR design for CoVs and would increase the options of molecular diagnosis for
56 circulating and emerging CoV infections. In this study, we designed and evaluated novel real-
57 time RT-PCR assays with locked nucleic acid (LNA) probes for clinically important CoVs based
58 on the identification of the abundantly expressed leader sequence in the 5'-untranslated region
59 (UTR) in small-RNA-Seq data analysis.

60 We included MERS-CoV (strain HCoV-EMC/2012, passage 8, provided by Ron
61 Fouchier, Erasmus Medical Center), HCoV-229E, HCoV-OC43, HCoV-NL63, and HCoV-
62 HKU1 in the study. SARS-CoV was not included as there has not been any human case since
63 2005. The MERS-CoV isolate was amplified by one additional passage in Vero cells to make
64 working stocks of the virus (5.62×10^5 50% tissue culture infective doses [TCID₅₀]/ml) as
65 previously described (18). All experimental protocols involving live MERS-CoV followed the
66 approved standard operating procedures of the biosafety level 3 facility at Department of
67 Microbiology, The University of Hong Kong, as previously described (19). High-titer stocks of
68 HCoV-229E, HCoV-OC43, and other respiratory viruses were prepared and their TCID₅₀ values
69 were determined using standard methods and as previously described (20,21-23). Attempts to
70 culture HCoV-NL63 and HCoV-HKU1 were unsuccessful because of their difficulty to grow in
71 cell lines available in our laboratories. Virus-positive clinical specimens (n=14) and laboratory
72 strains (n=13) used for evaluating the novel assays' cross-reactivities with other respiratory
73 viruses were obtained from archived nasopharyngeal aspirates at the clinical microbiology
74 laboratory at Queen Mary Hospital, Hong Kong. Total nucleic acid extractions of clinical
75 specimens and laboratory cell culture with virus strains were performed on 200 µl of sample

76 using EZ1 virus Mini Kit v2.0 (QIAgen) according to the manufacturer's instructions. The
77 elution volume was 60 μ l. Extracts were stored at -70°C or below until use. Total nucleic acid
78 extracts of ResPlex-II-HCoV-positive (n=49) and -negative (n=180) respiratory clinical
79 specimens prepared by using the QIAamp MinElute Virus Spin Kit were provided by the Hong
80 Kong Sanatorium and Hospital. A total of 229 fresh or frozen nasopharyngeal aspirates (NPAs)
81 collected between 1 January 2012 and 31 October 2014 from 229 pediatric and adult patients,
82 including 128 males and 101 females, aged 1 to 97 years, who were managed in Queen Mary
83 Hospital and Hong Kong Sanatorium and Hospital for upper and/or lower respiratory symptoms
84 were included in the study.

85 The most abundantly expressed sequence in the MERS-CoV genome was determined by
86 small-RNA-Seq data analysis (Supplementary Information). Approximately 2.6% of the trimmed
87 reads could be mapped onto the MERS-CoV genome. Among the mapped sequences, the
88 mapping analysis revealed that most of these small RNA sequence reads, accounting for >6,000
89 sequences (6.3%), matched the 67-nucleotide leader sequence at the 5' terminus of the genome
90 (Fig. 1). In contrast, the other peaks at the ORF1a, S, and N gene regions accounted for <3.0% of
91 the mapped small RNA sequence reads. Our mapping analysis also showed that the percentages
92 of mapped small RNA sequence reads at the gene regions targeted by the previously described
93 upE, ORF1a, ORF1b, N2, N3, NSeq, and RdRpSeq assays, which had longer sequences than the
94 67-nucleotide MERS-CoV leader sequence, were only 0.2%, 0.1%, <0.1%, 0.3%, 0.1%, 2.8%,
95 and 0.2% respectively (13,14,24). HCoV-229E, HCoV-OC43, HCoV-NL63, and HCoV-HKU1
96 similarly possess 70 to 72-nucleotide leader sequences at the same region in their respective
97 genomes (Fig. 1) (25-28). Although leader sequences of around 60 to 90 nucleotides in length
98 are found at the 5'-UTR upstream to the transcription regulatory sequence in the genomes and at

99 the subgenomic RNAs of all CoVs, the function of these leader sequences remains poorly
100 understood (29-31). In view of the abundance of the leader sequences and since infected cells are
101 known to contain large amounts of viral subgenomic RNA at which the leader sequences are
102 abundantly found (32,33), we hypothesized that the leader sequence might be a valuable
103 diagnostic target not only for MERS-CoV but also for other currently circulating HCoVs (Fig.
104 1).

105 To overcome the relatively short length of the leader sequences, we employed LNA
106 probes (Exiqon, Copenhagen, Denmark) to develop novel real-time RT-PCR assays for these
107 human pathogenic CoVs. LNA is a nucleic acid analogue with an extra bridge connecting the 2'
108 oxygen and 4' carbon that has exceptionally high hybridization affinity towards complementary
109 DNA and RNA and efficient mismatch discrimination (34). These properties are associated with
110 an increased melting temperature of the oligonucleotides, which allows the application of shorter
111 probes when LNA rather than DNA nucleotides are used in the nucleic acid amplification assays
112 (34). In recent years, LNA probes have been increasingly used in the design of real-time PCR
113 assays for other respiratory infections such as those caused by avian influenza A/H5N1 virus,
114 rhinovirus, enteroviruses, respiratory syncytial virus, and *Mycoplasma pneumoniae* (35-38).
115 Using LNA probes, we developed five novel real-time RT-PCR assays (named MERS-CoV-LS,
116 HCoV-229E-LS, HCoV-OC43-LS, HCoV-NL63-LS, and HCoV-HKU1-LS) targeting the short
117 leader sequences of these CoVs (Table 1).

118 The analytical sensitivities and specificities of the assays were excellent. The limits of
119 detection with *in vitro* RNA transcripts for MERS-CoV-LS, HCoV-229E-LS, and HCoV-OC43-
120 LS were 10 RNA copies/reaction and those for HCoV-NL63-LS and HCoV-HKU1-LS were 5
121 RNA copies/reaction. Linear amplification was achieved over an 8-log dynamic range, from 10^1 -

10⁸ RNA copies/reaction for all five assays, with calculated linear correlation coefficients (R^2) of 0.99-1.00 and amplification efficiency values of 1.93-2.27. The limits of detection with viral RNA were approximately 5.62×10^{-2} TCID₅₀/ml, 5.00×10^{-2} TCID₅₀/ml, and 3.16×10^{-3} TCID₅₀/ml for MERS-CoV-LS, HCoV-229E-LS, and HCoV-OC43-LS respectively (Supplementary Tables 1 and 2). The limit of detection for the MERS-CoV-LS assay was about one log TCID₅₀/ml higher than that for the MERS-CoV-upE assay in parallel runs and was comparable with those for the other assays currently recommended for screening and/or confirmation of MERS by the WHO, including the ORF1a, ORF1b, RdRpSeq, and NSeq assays (Supplementary Table 1) (7,14). Comparatively, the ORF1b assay for MERS-CoV has the least optimal limit of detection of 64 RNA copies/reaction (13,14). Our assays showed no cross-reactivity among the individual CoVs and with other common respiratory viruses including adenovirus, influenza A and B viruses, parainfluenza virus types 1 to 4, rhinovirus, respiratory syncytial virus, and human metapneumovirus (Supplementary Table 3).

Additionally, we assessed the diagnostic performance of our assays and compared it with ResPlex-II in in-use evaluation using 229 NPAs. ResPlex-II is a commercially available multiplex PCR assay which detects 18 respiratory viruses including HCoV-229E, HCoV-OC43, HCoV-NL63, and HCoV-HKU1 in a single run. It is commonly employed for laboratory diagnosis of viral respiratory tract infections in many clinical laboratories worldwide (39,40). Forty-nine NPAs which were tested positive for HCoVs by ResPlex-II and another 180 NPAs which were tested negative for respiratory viruses by ResPlex-II (Table 2) were tested in an operator-blinded manner. Our assays tested positive for the corresponding HCoVs with viral loads of 1.37×10^1 - 3.86×10^8 RNA copies/reaction in all 49 (100%) ResPlex-II-HCoV-positive NPAs (Table 2) ($p = 1.00$; Fisher's exact test). Moreover, our assays detected HCoVs in an

145 additional 4/180 (2.2%) NPAs which were initially tested negative by ResPlex-II, probably
146 because of the low viral loads of 2.29×10^1 - 2.40×10^2 RNA copies/reaction ($p = 0.12$; Fisher's
147 exact test). Sequencing analysis and two-step confirmatory real-time RT-PCR assays using
148 specific primers targeting the N gene of HCoV-OC43 and HCoV-NL63 (Supplementary
149 Information) confirmed that the results of these four ResPlex-II-HCoV-negative specimens
150 concurred with their CoV real-time RT-PCR assay results (two were positive for HCoV-OC43
151 and two were positive for HCoV-NL63). Overall, these results suggest that our assays are highly
152 sensitive and specific, and not inferior to ResPlex-II for the detection of HCoVs *in vitro* and in
153 clinical samples. It is important to note that while ResPlex-II and other multiplex PCR assays
154 have the advantage of being able to detect multiple viruses simultaneously, the sensitivity may be
155 <50% and inferior to monoplex PCR assays for HCoVs and other respiratory viruses such as
156 influenza A viruses (39,41). This relatively poorer sensitivity would especially limit the
157 application of these multiplex PCR assays for the detection of future emerging CoVs and avian
158 influenza A viruses which are potential pandemic agents that have significant public health
159 impact if a case was misdiagnosed.

160 Our study has demonstrated the previously unknown diagnostic value of the CoV leader
161 sequence and the usefulness of small-RNA-Seq data analysis in the selection of optimal gene
162 targets for the development of molecular diagnostic assays. The application of LNA probes
163 allowed the use of relatively short sequences such as the leader sequence of CoV genomes as a
164 diagnostic target in RT-PCR assays. The same approach may be applied to identify and design
165 real-time RT-PCR assays for other emerging viruses including novel CoVs that are likely to
166 emerge in the future, once their genomic data become available. As for any other gene targets
167 used in RT-PCR assays, particular attention should be paid to the presence of polymorphisms in

168 the leader sequences, which may affect the sensitivity of the assays. The novel CoV real-time
169 RT-PCR assays with LNA probes described in the present study should be further evaluated in
170 large-scale in-field evaluations. Development of these assays into multiplex assays with
171 comparable sensitivity and specificity and additional detection of other novel or re-emerging
172 CoVs may further enhance their clinical utility.

173

174 **ACKNOWLEDGEMENTS**

175 We thank Johnson Lau and Manson Fok for their facilitation of the study. This work is partly
176 supported by the donations of Mr. Larry Chi-Kin Yung, and Hui Hoy and Chow Sin Lan Charity
177 Fund Limited, the Consultancy Service for Enhancing Laboratory Surveillance of Emerging
178 Infectious Disease of the Department of Health, Hong Kong Special Administrative Region,
179 China, the University Development Fund and the Committee for Research and Conference
180 Grant, The University of Hong Kong.

181

182 **Ethical approval.** The study was approved by the Institutional Review Board of The University
183 of Hong Kong / Hospital Authority Hong Kong West Cluster.

184

185 **Conflict of interests:** None.

186

187 **FIGURE LEGENDS**

188 **Fig. 1** Schematic diagram of the MERS-CoV genome, with the leader sequence at the 5'-
189 untranslated region enlarged to illustrate the abundance of the small RNA sequences. The
190 percentages of mapped small RNA sequence reads at the leader sequence, ORF1a, S, and N gene
191 regions are quantified and shown. Leader sequences of 70 to 72 nucleotides in length are also
192 present in other human coronaviruses (HCoV-229E, HCoV-OC43, HCoV-NL63, and HCoV-
193 HKU1).

194 **TABLE 1** Primer and probe sequences of CoV real-time RT-PCR assays with LNA probes in the present study^a
 195

Assay	Genome target	Genome location	Primer/probe	Sequence (5' to 3')	GenBank accession no.
MERS-CoV-LS	Leader sequence	14-32	Forward	AGCTTGGCTATCTCACTTC	JX869059.2
		47-69	Reverse	AGTTCGTTAAAATCAAAGTTCTG	
		34-47	Probe	C+CT+CGT+T+CT+CT+TGC	
HCoV-229E-LS	Leader sequence	20-41	Forward	CTACAGATAGAAAAGTTGCTTT	NC_002645.1
		57-75	Reverse	ggTCGTTTAGTTGAGAAAAGT	
		44-59	Probe	AGACT+T+TG+TG+TCT+A+CT	
HCoV-OC43-LS	Leader sequence	17-28	Forward	aaaCGTGCGTGCATC	NC_005147.1
		43-66	Reverse	AGATTACAAAAAGATCTAACAAGA	
		32-48	Probe	C+TTCA+CTG+ATCT+C+T+TGT	
HCoV-NL63-LS	Leader sequence	23-46	Forward	ggAGATAGAGAATTTTCTTATTTAGA	NC_005831.2
		60-77	Reverse	ggTTTCGTTTAGTTGAGAAG	
		50-66	Probe	TGTGT+C+TAC+T+C+TTCT+CA	
HCoV-HKU1-LS	Leader sequence	21-37	Forward	CGTACCGTCTATCAGCT	NC_006577.2
		48-71	Reverse	GTTTAGATTTAATGAGATCTGACA	
		39-52	Probe	ACGA+T+CT+C+TTG+T+CA	

196 Abbreviations: HCoV, human coronavirus; MERS-CoV, Middle East respiratory syndrome coronavirus; UTR, untranslated region.

197
 198 ^a Probes were labeled at the 5' end with the reporter molecule 6-carboxyfluorescein (6-FAM) and at the 3' end with Iowa Black FQ
 199 (Integrated DNA Technologies, Inc). Lowercase letters represent the additional bases added which is not from the original genome
 200 sequence. The letters following "+" represent LNA bases which are modified with an extra bridge connecting the 2' oxygen and 4'
 201 carbon. The bridge "locks" the ribose in the 3'-endo (North) conformations and significantly increases the hybridization properties of
 202 the probe.

203 **TABLE 2** Comparison between CoV real-time RT-PCR assays with LNA probes and ResPlex-II for the detection of CoVs in
 204 nasopharyngeal aspirates
 205

Assay	Resplex-II-HCoV-positive NPAs ^a (n = 49)				Resplex-II-HCoV-negative NPAs ^a (n = 180)
	HCoV-229E	HCoV-OC43	HCoV-NL63	HCoV-HKU1	
MERS-CoV-LS	0/1	0/17	0/27	0/4	0/180
HCoV-229E-LS	1/1 (1.64×10^4)	0/17	0/27	0/4	0/180
HCoV-OC43-LS	0/1	17/17 (1.37×10^1 - 3.86×10^8)	0/27	0/4	2/180 (1.84×10^2 - 2.40×10^2)
HCoV-NL63-LS	0/1	0/17	27/27 (9.20×10^2 - 3.47×10^7)	0/4	2/180 (2.29×10^1 - 9.34×10^1)
HCoV-HKU1-LS	0/1	0/17	0/27	4/4 (1.94×10^3 - 4.42×10^5)	0/180

206 Abbreviations: NPAs, nasopharyngeal aspirates

207 ^a Values in brackets represent the ranges of quantitative results of positive samples in RNA copies/reaction.

208 REFERENCES

- 209 1. **Chan JF, To KK, Tse H, Jin DY, Yuen KY.** 2013. Interspecies transmission and emergence
210 of novel viruses: lessons from bats and birds. *Trends Microbiol* **21**:544-555.
- 211 2. **Chan JF, To KK, Chen H, Yuen KY.** 2015. Cross-species transmission and emergence of
212 novel viruses from birds. *Curr Opin Virol* **10**:63-69.
- 213 3. **Zaki AM, van Boheemen S, Bestebroer TM, Osterhaus AD, Fouchier RA.** 2012. Isolation
214 of a novel coronavirus from a man with pneumonia in Saudi Arabia. *N Engl J Med* **367**:1814-
215 1820.
- 216 4. **Cheng VC, Lau SK, Woo PC, Yuen KY.** 2007. Severe acute respiratory syndrome
217 coronavirus as an agent of emerging and reemerging infection. *Clin Microbiol Rev* **20**:660-694.
- 218 5. **Chan JF, Li KS, To KK, Cheng VC, Chen H, Yuen KY.** 2012. Is the discovery of the novel
219 human betacoronavirus 2c EMC/2012 (HCoV-EMC) the beginning of another SARS-like
220 pandemic? *J Infect* **65**:477-489.
- 221 6. **Chan JF, Lau SK, Woo PC.** 2013. The emerging novel Middle East respiratory syndrome
222 coronavirus: the "knowns" and "unknowns". *J Formos Med Assoc* **112**:372-381.
- 223 7. **Chan JF, Lau SK, To KK, Cheng VC, Woo PC, Yuen KY.** 2015. Middle East Respiratory
224 Syndrome Coronavirus: Another Zoonotic Betacoronavirus Causing SARS-Like Disease. *Clin*
225 *Microbiol Rev* **28**:465-522.
- 226 8. **Chan JF, Chan KH, Choi GK, To KK, Tse H, Cai JP, Yeung ML, Cheng VC, Chen H,**
227 **Che XY, Lau SK, Woo PC, Yuen KY.** 2013. Differential cell line susceptibility to the emerging
228 novel human betacoronavirus 2c EMC/2012: implications for disease pathogenesis and clinical
229 manifestation. *J Infect Dis* **207**:1743-1752.
- 230 9. **Muller MA, Raj VS, Muth D, Meyer B, Kallies S, Smits SL, Wollny R, Bestebroer TM,**

- 231 **Specht S, Suliman T, Zimmermann K, Binger T, Eckerle I, Tschapka M, Zaki AM,**
232 **Osterhaus AD, Fouchier RA, Haagmans BL, Drosten C.** 2012. Human coronavirus EMC does
233 not require the SARS-coronavirus receptor and maintains broad replicative capability in
234 mammalian cell lines. *mBio* **3**. pii: e00515-12.
- 235 10. **Chan KH, Chan JF, Tse H, Chen H, Lau CC, Cai JP, Tsang AK, Xiao X, To KK, Lau**
236 **SK, Woo PC, Zheng BJ, Wang M, Yuen KY.** 2013. Cross-reactive antibodies in convalescent
237 SARS patients' sera against the emerging novel human coronavirus EMC (2012) by both
238 immunofluorescent and neutralizing antibody tests. *J Infect* **67**:130-140.
- 239 11. **Lau SK, Che XY, Woo PC, Wong BH, Cheng VC, Woo GK, Hung IF, Poon RW, Chan**
240 **KH, Peiris JS, Yuen KY.** 2005. SARS coronavirus detection methods. *Emerg Infect Dis*
241 **11**:1108-1111.
- 242 12. **Song D, Ha G, Serhan W, Eltahir Y, Yusof M, Hashem F, Elsayed E, Marzoug B,**
243 **Abdelazim A, Al Muhairi S.** 2015. Development and validation of a rapid
244 immunochromatographic assay for the detection of Middle East Respiratory Syndrome
245 Coronavirus antigen in dromedary camels. *J Clin Microbiol* **53**:1178-1182.
- 246 13. **Corman VM, Eckerle I, Bleicker T, Zaki A, Landt O, Eschbach-Bludau M, van**
247 **Boheemen S, Gopal R, Ballhause M, Bestebroer TM, Muth D, Müller MA, Drexler JF,**
248 **Zambon M, Osterhaus AD, Fouchier RM, Drosten C.** 2012. Detection of a novel human
249 coronavirus by real-time reverse-transcription polymerase chain reaction. *Euro Surveill* **17**. pii:
250 20285.
- 251 14. **Corman VM, Muller MA, Costabel U, Timm J, Binger T, Meyer B, Kreher P, Lattwein**
252 **E, Eschbach-Bludau M, Nitsche A, Bleicker T, Landt O, Schweiger B, Drexler JF,**
253 **Osterhaus AD, Haagmans BL, Dittmer U, Bonin F, Wolff T, Drosten C.** 2012. Assays for

- laboratory confirmation of novel human coronavirus (hCoV-EMC) infections. *Euro Surveill* **17**. pii: 20334.
15. **de Sousa R, Reusken C, Koopmans M.** 2014. MERS coronavirus: data gaps for laboratory preparedness. *J Clin Virol* **59**:4-11.
16. **Sridhar S, To KK, Chan JF, Lau SK, Woo PC, Yuen KY.** 2015. A Systematic Approach to Novel Virus Discovery in Emerging Infectious Disease Outbreaks. *J Mol Diagn* pii: S1525-1578(15)00038-0.
17. **Corman VM, Olschlager S, Wendtner CM, Drexler JF, Hess M, Drosten C.** 2014. Performance and clinical validation of the RealStar MERS-CoV Kit for detection of Middle East respiratory syndrome coronavirus RNA. *J Clin Virol* **60**:168-171.
18. **Chan JF, Chan KH, Kao RY, To KK, Zheng BJ, Li CP, Li PT, Dai J, Mok FK, Chen H, Hayden FG, Yuen KY.** 2013. Broad-spectrum antivirals for the emerging Middle East respiratory syndrome coronavirus. *J Infect* **67**:606-616.
19. **Zheng BJ, Chan KW, Lin YP, Zhao GY, Chan C, Zhang HJ, Chen HL, Wong SS, Lau SK, Woo PC, Chan KH, Jin DY, Yuen KY.** 2008. Delayed antiviral plus immunomodulator treatment still reduces mortality in mice infected by high inoculum of influenza A/H5N1 virus. *Proc Natl Acad Sci U S A* **105**:8091-8096.
20. **Che XY, Qiu LW, Liao ZY, Wang YD, Wen K, Pan YX, Hao W, Mei YB, Cheng VC, Yuen KY.** 2005. Antigenic cross-reactivity between severe acute respiratory syndrome-associated coronavirus and human coronaviruses 229E and OC43. *J Infect Dis* **191**:2033-2037.
21. **Li IW, Chan KH, To KW, Wong SS, Ho PL, Lau SK, Woo PC, Tsoi HW, Chan JF, Cheng VC, Zheng BJ, Chen H, Yuen KY.** 2009. Differential susceptibility of different cell lines to swine-origin influenza A H1N1, seasonal human influenza A H1N1, and avian influenza

- 277 A H5N1 viruses. *J Clin Virol* **46**:325-330.
- 278 22. **Chan KH, Yan MK, To KK, Lau SK, Woo PC, Cheng VC, Li WS, Chan JF, Tse H, Yuen**
279 **KY.** 2013. Use of the human colorectal adenocarcinoma (Caco-2) cell line for isolating
280 respiratory viruses from nasopharyngeal aspirates. *J Med Virol* **85**:874-879.
- 281 23. **Li IW, To KK, Tang BS, Chan KH, Hui CK, Cheng VC, Yuen KY.** 2008. Human
282 metapneumovirus infection in an immunocompetent adult presenting as mononucleosis-like
283 illness. *J Infect* **56**:389-392.
- 284 24. **Lu X, Whitaker B, Sakthivel SK, Kamili S, Rose LE, Lowe L, Mohareb E, Ellassal EM,**
285 **Al-sanouri T, Haddadin A, Erdman DD.** 2014. Real-time reverse transcription-PCR assay
286 panel for Middle East respiratory syndrome coronavirus. *J Clin Microbiol* **52**:67-75.
- 287 25. **Herold J, Raabe T, Siddell S.** 1993. Molecular analysis of the human coronavirus (strain
288 229E) genome. *Arch Virol Suppl* **7**:63-74.
- 289 26. **Pyrce K, Jebbink MF, Berkhout B, van der Hoek L.** 2004. Genome structure and
290 transcriptional regulation of human coronavirus NL63. *Virol J* **1**:7.
- 291 27. **Woo PC, Lau SK, Chu CM, Chan KH, Tsoi HW, Huang Y, Wong BH, Poon RW, Cai JJ,**
292 **Luk WK, Poon LL, Wong SS, Guan Y, Peiris JS, Yuen KY.** 2005. Characterization and
293 complete genome sequence of a novel coronavirus, coronavirus HKU1, from patients with
294 pneumonia. *J Virol* **79**:884-895.
- 295 28. **Lau SK, Lee P, Tsang AK, Yip CC, Tse H, Lee RA, So LY, Lau YL, Chan KH, Woo PC,**
296 **Yuen KY.** 2011. Molecular epidemiology of human coronavirus OC43 reveals evolution of
297 different genotypes over time and recent emergence of a novel genotype due to natural
298 recombination. *J Virol* **85**:11325-11337.
- 299 29. **van Boheemen S, de Graaf M, Lauber C, Bestebroer TM, Raj VS, Zaki AM, Osterhaus**

- 300 **AD, Haagmans BL, Gorbalenya AE, Snijder EJ, Fouchier RA.** 2012. Genomic
301 characterization of a newly discovered coronavirus associated with acute respiratory distress
302 syndrome in humans. *mBio* **3**. pii: e00473-12.
- 303 30. **Qian Z, Dominguez SR, Holmes KV.** 2013. Role of the spike glycoprotein of human
304 Middle East respiratory syndrome coronavirus (MERS-CoV) in virus entry and syncytia
305 formation. *PLoS One* **8**:e76469.
- 306 31. **Yang D, Leibowitz JL.** 2015. The structure and functions of coronavirus genomic 3' and 5'
307 ends. *Virus Res* pii: S0168-1702(15)00115-X.
- 308 32. **Hui RK, Zeng F, Chan CM, Yuen KY, Peiris JS, Leung FC.** 2004. Reverse transcriptase
309 PCR diagnostic assay for the coronavirus associated with severe acute respiratory syndrome. *J*
310 *Clin Microbiol* **42**:1994-1999.
- 311 33. **Wu HY, Brian DA.** 2010. Subgenomic messenger RNA amplification in coronaviruses. *Proc*
312 *Natl Acad Sci U S A* **107**:12257-12262.
- 313 34. **Petersen M, Wengel J.** 2003. LNA: a versatile tool for therapeutics and genomics. *Trends*
314 *Biotechnol* **21**:74-81.
- 315 35. **Thanh TT, Pawestri HA, Ngoc NM, Hien VM, Syahrial H, Trung NV, van Doorn RH,**
316 **Wertheim HF, Chau NV, Ha do Q, Farrar JJ, Hien TT, Sedyaningsih ER, de Jong MD.**
317 2010. A real-time RT-PCR for detection of clade 1 and 2 H5N1 influenza A virus using locked
318 nucleic acid (LNA) TaqMan probes. *Virol J* **7**:46.
- 319 36. **Osterback R, Tevaluoto T, Ylinen T, Peltola V, Susi P, Hyypiä T, Waris M.** 2013.
320 Simultaneous detection and differentiation of human rhino- and enteroviruses in clinical
321 specimens by real-time PCR with locked nucleic Acid probes. *J Clin Microbiol* **51**:3960-3967.
- 322 37. **Do LA, van Doorn HR, Bryant JE, Nghiem MN, Nguyen Van VC, Vo CK, Nguyen MD,**

- 323 **Tran TH, Farrar J, de Jong MD.** 2012. A sensitive real-time PCR for detection and
324 subgrouping of human respiratory syncytial virus. *J Virol Methods* **179**:250-255.
- 325 **38. Komatsu H, Tsunoda T, Inui A, Sogo T, Fujisawa T, Imura M, Tateno A.** 2013.
326 Successful use of saliva without DNA extraction for detection of macrolide-resistant
327 *Mycoplasma pneumoniae* DNA in children using LNA probe-based real-time PCR. *J Infect*
328 *Chemother* **19**:1087-1092.
- 329 **39. Loens K, van Loon AM, Coenjaerts F, van Aarle Y, Goossens H, Wallace P, Claas EJ,**
330 **Ieven M; GRACE Study Group.** 2012. Performance of different mono- and multiplex nucleic
331 acid amplification tests on a multipathogen external quality assessment panel. *J Clin Microbiol*
332 **50**:977-987.
- 333 **40. Balada-Llasat JM, LaRue H, Kelly C, Rigali L, Pancholi P.** 2011. Evaluation of
334 commercial ResPlex II v2.0, MultiCode-PLx, and xTAG respiratory viral panels for the
335 diagnosis of respiratory viral infections in adults. *J Clin Virol* **50**:42-5.
- 336 **41. Chan KH, To KK, Chan JF, Li CP, Chen H, Yuen KY.** 2013. Analytical sensitivity of
337 seven point-of-care influenza virus detection tests and two molecular tests for detection of avian
338 origin H7N9 and swine origin H3N2 variant influenza A viruses. *J Clin Microbiol* **51**:3160-3161.

